

Use in medical offices, kitchens, and bathrooms on hard nonporous surfaces such as glass, stainless steel, Formica and other similar hard surfaces.

****KILLS:** Virucidal*, Bactericidal, Tuberculocidal, Fungicidal, Pseudomonacidal SaniPure Disinfecting Wipes are effective against the following pathogens in 2 minutes at 20°C/68°F, on hard, nonporous surfaces when used as directed:

Escherichia coli O157: H7, Methicillin-resistant Staphylococcus aureus (MRSA), Mycobacterium tuberculosis, Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Vancomycin-resistant Enterococcus faecalis (VRE), *Herpes Simplex Virus 2 (genital herpes virus), *Human Hepatitis B Virus, *Human Hepatitis C Virus, *Influenza A virus (A2 Japan). **Kills Mycobacterium tuberculosis.** SaniPure Disinfecting wipes are effective against the following pathogens in 5 minutes at 20 C/68 F, on hard, non-porous surfaces when used as directed. Trichopyton mentagrophytes (Athlete's foot fungus).

PREPARATION OF DISPENSER: Remove cover from canister and discard covering seal. From center of wipes roll, pull up a wipe corner, twist into a point and thread through the tear throat in the underside of the lid cover. Pull wipe through about one (1) inch. Replace the lid cover on container, assuring a secure fit. Pull out first wipe and snap at a 90° angle. Remaining wipes feed automatically for next use. Always reseal hinged cap after each use to ensure moisture retention.

DIRECTIONS FOR USE: It is a violation of U.S. Federal law to use this product in a manner inconsistent with its labeling. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection. This product is not for use on the inhalation therapy and/or pulmonary diagnostic equipment.

DISINFECTION: Thoroughly wet pre-cleaned, hard, nonporous surface with a wipe, keep wet for 2 minutes (5 minutes if fungus is suspected), and allow to air dry. Use as many wipes as needed for the treated surface to remain wet for the entire contact time. For use as a general disinfectant wipe on hard, nonporous, inanimate surfaces. You may use a wipe to pre-clean the surface to be disinfected, but thorough re-wiping with a fresh wipe is required for disinfection. Use on food-contact surfaces must be followed by a potable water rinse.

KILLS VIRUSES* & BACTERIA**

For use on environmental surfaces

**Details on back



Disinfecting Wipes

ACTIVE INGREDIENTS:

N-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
Dimethyl Benzyl Ammonium Chloride.....	0.12%
N-Alkyl (68% C12, 32% C14) Dimethyl Ethyl	
Benzyl Ammonium Chloride.....	0.12%
Isopropyl Alcohol.....	41.58%
OTHER INGREDIENTS†	58.18%
TOTAL	100.00%

† Does not include weight of dry wipe.

KEEP OUT OF REACH OF CHILDREN
WARNING:
DO NOT USE AS A BABY WIPE

See side panels for additional precautionary statements.



160 Presaturated Wipes

NET CONTENTS: 1lb. 10 oz.
(6 in. by 6.7 in.) (15cm x 17cm)

STORAGE AND DISPOSAL:

Store in cool area. Keep away from heat or open flame. Do not reuse wipes. Dispose of used wipe in trash. Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

PRECAUTIONARY STATEMENTS:

Hazards to Humans and Domestic Animals

WARNING: Causes substantial, but temporary eye injury. Do not get in eyes or on clothing. Wear specific, appropriate protective eyewear such as goggles, face shield or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear disposable latex gloves, gowns, masks and eye coverings when handling HIV-1 infected blood or body fluids. Avoid contamination of food. Remove and wash contaminated clothing before reuse.

FIRST AID:

If in eyes: Hold eye open and rinse slowly and gently with water for at least 15–20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice. Wash contaminated clothing before reuse. **WASH THOROUGHLY AFTER HANDLING WIPES.**

Emergency Number: 1-800-222-1222

Manufactured for:

Rx SAN, LLC
1877 E. 17th Ave.
Columbus, OH 43219



BATCH NUMBER:



EPA No.: 59894-10-99867

EPA Establishment No.: 52823-OH-1